

Title: Domestication as a model system for the Extended Evolutionary Synthesis

Author: Melinda A. Zeder

Affiliation 1: Program in Human Ecology and Archaeobiology, Department of Anthropology, National Museum of Natural History, Smithsonian Institution, 10th and Constitution, Washington DC, 20560, USA

Affiliation 2: Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA

Abstract: One of the key challenges in evaluating arguments for extending the conceptual framework of evolutionary biology involves the identification of a tractable model system that allows for an assessment of the core assumptions of the extended evolutionary synthesis (EES). The domestication of plants and animals by humans provides one such case study opportunity. Here I consider domestication as a model system for exploring major tenets of the EES by first discussing the novel insights that Niche Construction Theory (NCT, one of the pillars of the EES) provide into the domestication processes, particularly as they relate to five key areas: co-evolution, evolvability, ecological inheritance, cooperation, and the pace of evolutionary change. This discussion is next used to frame testable predictions about initial domestication of plants and animals that contrast to those grounded in standard evolutionary theory, and to demonstrate how these predictions might be tested in multiple regions where initial domestication took place. I then turn to a broader consideration of how domestication provides a model case study consideration of the different ways in which the core assumptions of the EES strengthen and expand our understanding of evolution, including reciprocal causation, developmental processes as drivers of evolutionary change, inclusive inheritance, and the tempo and rate of evolutionary change.

Key Words: Domestication, Niche Construction Theory, Co-Evolution, Evolvability, Ecological Inheritance, Cooperation

Review article:

Domestication as a model system for the extended evolutionary synthesis

Melinda A. Zeder

Interface Focus 2017 7 20160133; DOI: 10.1098/rsfs.2016.0133. Published 18 August 2017

<http://rsfs.royalsocietypublishing.org/content/7/5/20160133>

1. Introduction

First proposed in the mid-twentieth century, the Modern Synthesis (MS) has proven a powerful conceptual framework for studying evolution that extends far beyond biology into the social sciences and the humanities. By bringing Mendelian genetics together with neo-Darwinism, the MS provided the mechanisms on which Darwinian selection operated, and that, through the agency of population genetics, could be operationalized into a coherent understanding of the process of evolution [1-3]. In the MS, natural selection is recognized as the preeminent and ultimate causal force in evolution, with variation arising through random mutation, and adaptive variations passed on at a higher rate than less adaptive ones, resulting in an evolutionary process that proceeds at a gradual pace made up of small micro-evolutionary changes in the composition of individual genes and alleles within genes (Table 1).

Early calls for emendation of the core principles of the MS noted the importance of directional processes that sort and pass on variation in ways that may not always conform to adaptive fitness criteria, and envisioned evolution as proceeding at an uneven rate with long periods of stasis punctuated by events of major macro-evolutionary change [4-6]. Subsequent insights from developmental biology, ecology, and the social sciences have built on these initial calls for revision, highlighting areas in which core MS principles may require additional reevaluation and extension. These include insights about the evolutionary impacts of ecosystem engineering in influencing the selective environments of both engineers and others sharing constructed niches - and their descendants [7]; about the role of multiple constructive developmental processes in leading rather than following genetically-driven change [8]; and the recognition of additional inheritance systems - internal and external to an organism - that shape evolutionary trajectories [9,10]. These insights have recently been integrated into a general call for a revision of the classical MS by an Extended Evolutionary Synthesis (EES) [11,12]. Proponents of the EES maintain that, in addition to natural selection, other powerful forces shape evolution through interactive bouts of development and selection operating on variation that, along with genetic mutation, arises via the agency of constructive processes. They further contend that these variations are passed from one generation to the next by

multiple interacting inheritance systems, resulting in directional, non-random change at the level of genes, whole organisms, or even groups of organisms, and that evolution is more likely to proceed at variable, often quite rapid, rates (Table 1).

One of the major challenges in making the case for the EES is to demonstrate that the core principles of the EES can generate empirically testable predictions about evolutionary process that are not considered under the MS [13-15]. While many studies have attempted to empirically evaluate elements of the EES framework, advocates of emendation recognize that more is needed.

“Although many systems can reveal individual pieces of a larger puzzle, the gold standard is ... for a single system ... to understand the complete set of links between environmental variation, developmental responses, and changes to genetic and epigenetic architecture over the course of evolution [16:666].

Here I argue that the initial domestication of plants and animal by humans provides a compelling example of such a gold standard system. Domestication is a process with clear and dramatic evolutionary impact. It created new organisms that cannot live outside of a domesticatory relationship with humans; contributed to significant declines in overall biodiversity through the extirpation or even extinction of free-living species; transformed landscapes and atmospheric conditions; and reshaped the trajectory of human cultural evolution [17]. Beyond its unquestionable status as a major evolutionary event, what makes domestication an exceptionally valuable model system is that it encompasses all the central issues involved in ongoing debates over the need for an extended evolutionary synthesis – from the developmental processes EES advocates maintain require greater attention to the multiple forms of inheritance held to shape evolutionary trajectories. Moreover, new archaeological, archaeobiological, and genetic data are producing robust records of initial domestication in multiple world areas, all set within an increasingly well-resolved temporal framework that allows for an assessment of how these multiple constructive processes and inheritance systems affect the pace, tempo, and direction of evolutionary change.

The case for domestication as a model system for exploring major tenets of the EES is made here first by discussing the novel insights that Niche Construction Theory (NCT, one of the pillars of the EES) provide into the domestication processes as they relate to five key areas: co-evolution, evolvability, ecological inheritance, cooperation, and the pace of evolutionary change. This discussion is next used to frame testable predictions about the initial domestication of plants and animals that contrast to those grounded in standard evolutionary theory, and to demonstrate how these predictions might be tested in the many different world areas that witnessed the independent domestication of a wide range of different species of plants and animals. I conclude with a broader consideration of the value of domestication as a model system for evaluating core assumptions of the overall EES as it relates to issues of reciprocal causation, the role of constructive developmental processes, inclusive inheritance, the targets of selection, and rates of change.

2. Explanatory Frameworks for Initial Domestication

Archaeologists have a long history of looking to evolutionary biology as a source of explanatory frameworks for initial domestication. To date, most of these frameworks have been grounded in neo-Darwinian principles. One approach, promoted by researchers adhering to the selectionist school of archaeological theory, embraces a MS trait-level view of evolution (with individual human behaviors and their artifact proxies taking the place of genes and alleles) and envisions the process of cultural evolution as made up of small micro-evolutionary trait adjustments [18-22]. Another is based in optimal foraging theory, which is itself a theory borrowed from micro-economics that self-identifies with Neo-Darwinism on the assumption that foragers who optimize for returns enjoy selective advantage over those who do not [23-25]. Both selectionist and OFT explanations characterize domestication as a slow incremental co-evolutionary process between humans and target plant and animal species initiated in the context of resource depression, caused either by environmental change or by human impacts on resource availability, which forced foragers to turn to lower ranked, less preferred resources (such as the progenitors of domesticates) garnered through increasingly more intensive delayed-return strategies. Natural selection serves as the primary driving force, guiding both

the co-evolutionary relationships between humans and target species, as well as the individual decisions foragers make about resource choice and exploitation strategy (for a more extended discussion of neo-Darwinian explanatory frameworks for initial domestication see references 26 and 27).

More recently archaeologists have begun to look to niche construction theory (NCT), for an alternative explanatory framework for initial domestication [27-34]. NCT focuses on the evolutionary impacts of ecosystem engineering activities that modify selection pressures acting on present and future generations living within altered niches [7]. Broadly defined, NCT encompasses the niche-altering activities of all organisms, with humans characterized as “the ultimate niche constructors” [7:28]. The inclusion, and indeed emphasis, in NCT on “cultural niche construction” has drawn social scientists to NCT who see it as a way of bridging the divide that has traditionally separated the study of cultural evolution from that of biological evolution [35-40]. This has been particularly true of archaeologists interested in the origins of plant and animal domestication [27-34]. An obvious strength of NCT has been the extent to which it provides special insight into domestication in five key areas: co-evolution, evolvability, ecological inheritance, cooperation, and rates of change [33:Table 2].

2.1 Co-evolution

A central tenet of NCT is that niche-constructing activities cause eco-evolutionary feedbacks in which “ecological interactions drive evolutionary change in organismal traits that, in turn, alter the form of ecological interactions” [41:1629,42]. Eco-evolutionary feedbacks that arise in the context of niche construction are likely to foster co-evolutionary interactions between organisms living in constructed niches [43-45]. These relationships are considered diffuse when the impacts of the niche-constructing activities of one species on another are mediated by some change in abiotic conditions or intermediary biota. Diffuse co-evolutionary interactions may have wide ranging evolutionary and ecological effects that ripple outward from the niche-constructing organism and affect other organisms in ways that may or may not have a reciprocal impact on the original niche-constructor. Co-evolutionary relationships are particularly potent, however, when they are pairwise and involve interacting organisms that are

both actively engaged in and affected by niche-constructing activities that drive reciprocal evolutionary responses in each [45].

NCT insights into the development of co-evolutionary relationships in the context of niche-construction have clear and obvious implications for understanding domestication which is best defined as a pairwise, multi-generational co-evolutionary mutualism between a domesticator and a domesticate that arises in the context of niche-construction [32,33]. By assuming some degree of influence over the care, protection, and/or reproduction of the target domesticate, the domesticator engages in niche-constructing activities aimed at increasing the supply or predictability of a resource provided by the domesticate, while the domesticate, in turn, engages in reciprocal niche-constructing activities that enable it to benefit from its association with humans in a way that enhances its fitness over con-species that remain outside this relationship. And this is true of all the major pathways that humans and target species have followed into domestication [46]. The harvest, or prey, pathway, for example, is initiated by humans through eco-system engineering activities that manipulate the conditions of growth of an organism or its environment with the goal of increasing its relative abundance or predictability. Moving plants into protected areas, channeling water, landscape burning, building traps or corrals, selective culling can all be considered niche-constructing activities initiated by humans that can result in the development of co-evolutionary relationships between human niche-constructors and target plant or animal populations that may lead to domestication [29]. This is the pathway that brought many perennial plant and tree species, as well as most livestock animals, into domestication. The commensal pathway, on the other hand, is initiated by plants and animals that, through relocative niche-construction, move into anthropogenic environments to take advantage of new opportunities available there. Many “weedy” annual plants followed this pathway into domestication as they colonized soils disturbed by anthropogenic activities [47], as did animals such as wolves and wild cats that ventured into human settlements to feed off human refuse or to prey on other animals drawn to these environments [46]. Finally, the directed pathway to domestication in which humans deliberately set out to domesticate a species to extract some resource of interest also involves a significant degree of niche-construction as humans tightly control the target species’

movements, nutrition, and breeding (to the point of direct genetic manipulation). This is the pathway followed by all recent domesticates, with the explosion of aquatic species brought into domestication in the last 50-100 years [48] - whose rearing requires major transformation of aquatic environments – being a particularly compelling example of the role of niche-construction in directed domestication.

In each case, however, the co-evolutionary relationships that develop in these contexts will only result in domestication if the niche-constructing activities initiated by one partner (the domesticator or the domesticate) are met by those of the other in ways that enhance the benefits both derive from the relationship, increasing the mutual investment of each in maintaining it [32]. Domestication, then, requires a sustained multi-generational co-evolutionary relationship between two niche-constructing species with each undergoing changes (either genetically-driven or plastic) that enhance the benefits each derives from the relationship and/or that make further investment in the relationship more attractive to its partner. When humans sow seed in prepared substrates, for example, receptive plants may respond by altering the timing of germination and subsequent ripening - a form of developmental niche-construction [49] - in ways that increase the likelihood of their harvest by humans and subsequent inclusion in the next season's seed stock [50]. Animals practicing relocative niche construction by moving into anthropogenic environments experience selection for reduced reactivity to humans that enhances their reproductive success in that setting and encourages reciprocal niche-constructing activities on the part of humans directed toward the animals' care, protection, and breeding [46,51].

2.2 Evolvability

Whether organisms engage in co-evolutionary relationships in the context of niche construction depends in large measure on their evolutionary potential or evolvability – their ability to respond to the new selective pressures in constructed niches in ways that result in heritable evolutionary change in descendant populations [41,42]. Insights from NCT about the potential of organisms to respond to altered-niches in evolutionary significant ways have special resonance for understanding the process of domestication that merit a more extended consideration.

2.2.1 Evolutionary potential

In domestication, the capacity of different potential partners to respond to new selective pressures created by niche construction plays a major role in determining which will embark on pathways leading to domestication and the directions those pathways will take. In plants, annual species with fewer specific habitat requirements and which are able to colonize disturbed habitats are thought to be good candidates for domestication [47]. Behavioral traits such as reduced wariness, hierarchical herd structure, and generalized feeding and habitat preferences make certain animal species more likely potential domesticates than species that lack these traits [46]. Variables that make certain human groups more likely to engage in sustained co-evolutionary relationships leading to domestication include their willingness/ability to modify technology and alter existing subsistence strategies in light of new opportunities or challenges, their assessment of the current and potential demands and returns of available resources, as well as their physiological capacity to utilize resources [32,33]. In each case, on both the plant/animal and human side of the equation, a certain degree of plasticity or flexibility in the ability to take advantage of new opportunities and respond to new challenges is an important pre-requisite for embarking on pathways to domestication.

2.2.2 Phenotypic plasticity

To a major degree this flexibility may be attributable to a capacity for engaging in constructive developmental processes that shape evolutionary trajectories in non-random, directional ways [12,52]. Of these processes, the ability of an organism to match its phenotype to different environments through the plastic expression of novel morphological, physiological, or behavioral traits [53] may be the one of the most important factors behind an organisms' potential for engaging in a domesticatory relationship [54]. To date most of the research on the role of phenotypic plasticity in domestication has concentrated on the impact of climate change in the expression of cryptic variation in progenitors of crop plants. Experimental studies have, for example, demonstrated that teosinte grown under reduced CO₂ atmospheric conditions, similar to those of the Late Pleistocene/Early Holocene, exhibit a number of maize-like traits not seen in teosinte grown under modern CO₂ conditions [55]. These traits would have made

these plants more productive and easier to harvest, it is argued, leading to co-evolutionary relationships between foragers and certain teosinte morphotypes that, eventually, resulted in the fixation of these traits in domesticated maize. An additional trigger of plastic responses to Early Holocene climate change may be the impact of more marked, but predictable seasonality in climate during this time [56], environmental conditions which Donohue posits may be responsible for triggering “phenological cuing” in organisms - a form of developmental niche construction in which phenotypic responses to conditions experienced in one stage of an organism’s life cycle determine those displayed during subsequent life stages [49].

Another promising area for future research is the role of niche-construction in setting parameters for the accumulation and expression of phenotypic plasticity in emergent domesticates. Exposure to novel environments can result in the release of otherwise cryptic variation that facilitates rapid adaptive evolution in receptive species [57-59]. This process of developmental selection involves sampling a range of phenotypes, with feedback from the novel environment helping to reinforce the high performing phenotypes [60]. The introduction of plants and animals to novel environments created by anthropogenic disturbance or manipulation is then likely to have resulted in the expression of otherwise cryptic phenotypic variation in species with the ability to alter phenotypes in response to different environmental conditions, allowing plants and animals entering anthropogenic environment to rapidly adjust to the new selective pressures encountered there. Contemporary studies, in fact, find that plastically-induced trait changes occur in much higher rates in organisms entering anthropogenic environments than in natural contexts [61], with human harvesting of organisms being a particularly potent factor in inducing dramatic phenotypic changes [62-64].

Once within these environments, plants and animals living under human management and care would have been shielded from many of the selective pressures faced by con-species living outside this relationship, creating the kind of homeostatic mechanisms thought to encourage the accumulation of additional genetic variation and novel combinations of variants in populations that may be released at some later point [16,60]. The effect of human agency in buffering managed plants and animals may have played an especially important role in reducing the costs of developmental selection, which include expenditures of energy and material

needed to express different phenotypes and the risks incurred by the expression of potentially deleterious traits during the trial-and-error exploratory phase of adjustment to a new environment [53]. Human intervention and buffering, in fact, frequently favored traits in target species of interest to humans that would otherwise be selected against outside the managed environment (i.e., indehiscent seed dispersal traits that increase the yield of human harvested plants, but that face strong negative selective pressures in the wild [50]).

The release of cryptic variation within anthropogenic environments may well, then, be responsible for the high degree of variability in trait expression documented in crop plants during initial stages of domestication, as has been suggested for the non-directional expression of variable seed sizes and coat thickness in managed chenopodium in Eastern North America [65]. Similarly, the high degree of variability observed in the expression of the non-shattering rachis in cereals and in seed size among both cereals and pulses during initial stages of their cultivation in the Near East, commonly attributed to restocking or introgression of managed plants with free-living ones [66], may, instead, be attributable to the release of phenotypic plasticity in plants living within anthropogenic managed environments.

2.2.3 Pleiotropy

Plastic responses to novel environments may also elicit the expression of pleiotropically linked traits across the life cycle of an organism in ways that can both foster and constrain its potential for evolutionary change [49,57,60,67] – a process that may be responsible for the constellation of traits that often co-occur in domesticates. In plants [47,68] these include morphological and phenological traits that affect: 1) germination (changes in dormancy thresholds, changes in seed size and tests thickness), 2) dispersal (synchronization in ripening, development on non-shattering seed heads and pods), 3) plant architecture (reduction in number of branches, plant stature, apical dominance), and 4) productivity (increases in density of plants and in the number and size of plant organs of economic interest to humans). In animals [46] these include: 1) reduction in brain size (especially in regions of the brain that control aggression and reactivity to outside stimuli), 2) developmental rates and sequences (usually a delay in rates of development or paedomorphosis), 3) a corresponding neotonization in morphology (often

manifested in changes in cranial morphology), 4) retention of juvenile behaviors, 5) early sexual maturity, 6) lop ears (thought to be the result of a failure of cartilage to fully develop due to early sexual maturity), and 7) piebald coats (possibly an artifact of the shared pathways traveled by melatonin and serotonin, an example of the way in which similar signaling and resource mobilization pathways can induce pleiotropic responses in organisms [49]).

Some researchers maintain that these constellations of domestication traits are best viewed as pleiotropic cascades that arise from the orchestration of gene expression during development caused by mutations in a limited number of regulatory genes [69]. Earlier studies suggested that this was indeed the mechanism for the manifestation of these traits in domesticates. A comparative mapping of Qualitative Trait Loci in pearl millet, for example, suggested that the same loci controlling spikelet structure in this species were also responsible for seed shattering in maize, rice, and sorghum [70]. Another early study demonstrated the pleiotropic linkage between loci responsible for the expression of traits affecting growth habit and phenology, seed dispersal and dormancy, and fruit and seed size in the common bean [71].

Subsequent research employing whole genome sequencing and genome-wide association studies, however, has shown that the origins of these traits is more complicated than originally thought. The non-shattering trait found in domestic cereals, for example, is now known to be controlled by different genes in different species, with, in some cases, different linked loci implicated in the formation of the tough rachis in different independent domestication events involving the same species [50]. Studies of gene expression in different pairs of domesticated and wild animals found little or no overlap in genes or gene complexes controlling wariness and aggression, and no evidence for a genomic association between behavior and coat color in domesticated rats [51, 72, 73]. Another recent study suggests that linked traits in domestic animals arise not through the action of individual genes or gene complexes, but through the multi-genic down-stream consequences of selective factors operating on neural crest cells during embryonic development [74] - an example, perhaps, of the role that organismal levels above genes play in an organism's ability to shape its own developmental trajectory [12]. Another study maintains that neuro-endocrine mechanisms,

specifically the role of single hormones in targeting multiple tissues, underlie the expression of pleiotropic behavioral traits seen in domestic animals [75].

2.2.4 Hybridization

Another way in which novel traits can be acquired is by horizontal transfers of DNA through hybridization and genome doubling that can result in major epigenetic change in a single generation [76,77]. Introgressive hybridization and polyploidy is a common feature of many domestic crop species (i.e. cotton, bread wheat, tobacco, sugar cane, potatoes and other related tubers), with sometimes complicated histories of auto- and allo-polyploidy in the same species occurring both prior to and concurrent with human manipulation [78]. Hybrid origins without gene doubling has been suggested for at least one common animal domesticate, with the yellow skin characteristic of the domestic chicken shown to be the result of the hybridization of the red junglefowl with the closely related grey junglefowl [79]. And we know that at least the later history of South American camelids has been shaped by a complicated process of hybridization between domesticated llamas and alpacas and their progenitors, guanacos and vicuñas [80]. The capacity for successful hybridization resulting in dramatic genomic reorganization, or “cataclysmic evolution” [76,77], may, in fact, have been another trait that predisposed certain species to rapidly adapt to anthropogenic environments and eventual domestication, as well as one that assisted in the dispersal of domesticates [51,78].

2.2.5 Niche construction and constructive development

While the mechanics of the acquisition and expression of these domestication traits is an active and productive area of study, it may be equally interesting to focus on the role of niche construction in creating the selective environments in which they arose in the first place. Viewing these traits as constellations of environmentally mediated genetic associations (EMGAs in NCT parlance [44]) that arise in the context of altered environments raises a number of promising new avenues of inquiry. One direction for future research in this vein would be examining the role of co-constructed niches in generating selective pressures that triggered the appearance of the same constellation of traits in different species. Such research might

investigate the commonalities in human/plant interaction that resulted in the same constellation of traits in different cereal crops such as wheat, barley, rice, and millet domesticated (in some cases multiple times) in the Near East, East Asia, and Africa [50]. Another promising goal for future research would be identifying the conditions that trigger plastic responses and pleiotropic cascades in certain species within certain engineered environments, but not others living in the same or similar environments. Continuing with the example of cereals, here one might ask why wild oats that were intensively collected in the Early Holocene Levant failed to move on to domestic status at this time, while wheat and barley that were exploited along with wild oats responded to human manipulation by developing traits that led to their domestication [81]. A better understanding of how these co-constructed niches change selective pressures on plants and animals undergoing domestication may give new insight into the variability in the timing and sequence of the appearance of different domestication traits in different species. A goal might here be discovering why seed size increase preceded the development of indehiscence in Near Eastern cereals (emmer and einkorn wheat and barley) by hundreds of years or more, while in Near Eastern pulses (chickpeas, fava beans, and lentils) indehiscence seems to have developed long before any clear trends toward seed size increase, even though these crops were brought into domestication at the same time by the same early cultivator groups [50,82]. And finally, new research might explore how domesticatory relationships arising in the context of co-constructed niches served to buffer certain species from the deleterious effects of hybridization and encouraged the perpetuation of traits that outside this relationship would have conferred no selective advantage or even been selected against, but that within it proved highly advantageous to both human and the domesticate. A profitable line of inquiry here would be examining the role of humans in fostering crosses between a species of goat grass and domesticated emmer wheat – itself derived from a polyploid progenitor created by a cross between other species of goat grass and wheat – to form bread wheat, a hexaploid free-threshing and highly productive wheat species grown today around the world but not found in the wild [83].

2.2.6 Genetic accommodation and assimilation

In order for traits arising in the context of niche construction to have evolutionary impact there must be some mechanism by which they are stabilized within a population and become heritable traits with evolutionary impact - a process EES advocates call genetic accommodation [57-59,84,85]. In domesticates genetic accommodation of certain traits may come about through the adoption of new harvesting, culling, or processing practices, as has been proposed for the fixation of the non-shattering rachis in wheat and barley once sickles were used as the primary harvesting tool [86 but see 87]. Deliberate human selection for certain desirable traits may also have been responsible for their fixation, as may have been the case in the selection for the woolly undercoat, visible only during molting season in wild sheep, that became the dominant fiber in the coats of domestic sheep [88]. But perhaps the most significant factor in this process was the movement of managed plants and animals into new areas where factors like genetic drift and founders effect, responses to new selective pressures encountered in these new environments, and, especially, reproductive isolation from free-living sympatric served to fix traits that arose in the context of initial management.

Domesticates likely also underwent a process of genetic assimilation [16,53, 58,59,84,89], a sub-set of genetic accommodation, that results in the reduction or loss of plasticity, and the costly machinery underlying plasticity, so that the plastic traits that made the plant or animal an attractive domestic partner to humans no longer required environmental cuing for expression. Despite the loss of plasticity and the major genetic bottlenecks caused by factors such as genetic drift, founders effect and reproductive isolation [50], domesticates likely retained at least some capacity for the plastic expression of traits that that made it possible to successfully disperse far beyond the natural habitats of their progenitors and the homeland of initial domestication. The dispersal of Near Eastern cereal crops into Europe, for example, could only have been accomplished if these domesticates retained the ability to undergo changes in genetic controls on seasonality, especially in flowering phenology and accommodations for length of growing season, that allowed them to adjust to the very different seasonal regimes and climatic conditions they encountered in Europe [90,91]. Similarly, plasticity in the expression of vernalization genes among different land races of wheat and barley may have been key in determining the pace and direction of their eastward dispersal across and around

the Tibetan Plateau into Asia [90, 92], as well as the dispersal of rice out of the Yangtze basin into northern China [90], and of maize from central Mexico throughout North America [93-95].

2.3 Ecological inheritance

A basic tenet of NCT is that the niche-altering activities leave an ecological inheritance that persists beyond the lifetime of the original niche-constructing organism and bequeaths a legacy of modified selection pressures to its own and other organisms' descendants [96,97]. This concept has obvious relevance to the niche-constructing activities by humans and plants/animals involved in emergent domestication. Human landscape management aimed at promoting species of economic interest and their alterations of the population structure of biotic communities, as well as the relocative and developmental niche constructing activities of plants and animals living within anthropogenic environments, all serve to create reinforcing eco-evolutionary feedbacks that deepen the commitment of all partners to maintaining the co-constructed ecological niche that they bequeath to their descendants [33].

The impact of this ecological legacy in shaping the evolutionary trajectories of subsequent generations is so great, NCT advocates argue, that it qualifies as a second general inheritance system on a par with genetic inheritance [7, 97]. Further, they maintain that there are multiple interacting channels over which information is transmitted to subsequent generations [98,99]. Channels internal to an organism include not only genetic inheritance, but also developmental processes like epigenetic inheritance and maternal effects. External channels that operate outside the organism include the inheritance of modified selective environments and, importantly, acquired and learned behaviors. While template copying systems like genetic inheritance ensure a high degree of fidelity in information transfer over internal channels, the targets of these high-fidelity transfers are limited to parental organisms and their direct descendants. Information transferred through external channels, on the other hand, is more prone to transcription and other errors, but its targets are far wider [100]. One way to increase the fidelity of information transfer over external channels, at least in animals, is through improvements in the transmission of acquired behaviors that enhance an animal's ability to learn from one another and allow novel traits to spread widely through a population

[43]. In humans, the capacity for language and symbolic communication vastly enhances information transfer making it possible to modify and fine-tune behaviors and so “ratchet” up their complexity and efficiency [101].

Small-scale human societies, such as those engaged in initial domestication, use this capacity to amass stores of ecological knowledge (Tradition Ecological Knowledge or TEK) that become the primary vehicles for transmitting this ecological inheritance to their descendants [30,102]. Extensively documented in small-scale societies around the world [103-106], TEK consists of continually updated cognitive maps of resource distributions, information about the life cycles of economically important resources, their seasons of availability, behavior, and the ways in which environments or biotic communities can be manipulated to enhance the supply and predictability of target resources [30]. Played out in stories, myths, ritual performance, and lessons passed on from elders to younger members of social groups, TEK provides a coherent framework for how the world works and the human place within it [30,102]. Protecting, preserving, and continually updating this store of knowledge is essential for the survival of societies engaged in the kinds of ecosystem engineering that both forges and deepens the co-evolutionary dependencies between humans and emergent domesticates. Including these cultural behaviors among the external channels that shape evolutionary trajectories, then, provides fresh insight into the how the generation and transmission of TEK plays a catalytic role in the domestication process.

2.4 Cooperation

By modulating resource availability, niche-constructing activities create the potential for a tragedy of the commons in which some individuals reap the benefits of resource enhancing activities without contributing to them [107]. The perpetuation of constructed niches that form the basis of ecological inheritance, and the means by which niche construction becomes an evolutionary process, requires conditions that counter disincentives for cooperation and reward individuals that contribute to collective activities. Clearly there are pressures for cooperative behaviors among human groups involved in the niche-altering activities leading to

domestication, at the very least to combat the advantage that cheaters would enjoy from the collective activities that they themselves shirk [108].

One way in which cooperative behaviors can be encouraged, NCT advocates argue, is by broadening the focus of the rewards of collective actions from those enjoyed by an individual over contemporary competitors, to those that the individual bequeaths to relatives living in the future. In this way activities of the individual that benefit relatives living in the future can be seen as directly benefiting the individual [109]. This form of “trans-generational altruism” [110] is particularly effective in philopatric organisms whose long-term investments in altered niches make it more likely that they and their descendants will continue to reap the benefits of these collective activities. Incentives for cooperative behaviors, in contrast, are diminished under conditions of high mobility and dispersal where there is a smaller pay-off from niche-altering behaviors and a decreased probability of relatedness among aggregating individuals [109]. Another factor thought to encourage the cooperative behaviors is the ability to monopolize constructed niches thereby ensuring that niche-constructing populations continue to benefit from collective activities, while minimizing the extent to which interlopers are able to take advantage of them [107]. It is likely no accident, then, that initial domestication of plants and animals usually took place in the context of semi- to fully sedentary groups that had long term investments in the exploitation of abundant, diverse resources that could be predictably found within well-defined catchment areas [27,30,111]. These are contexts that could have been both monopolized and defended in ways that encouraged the kinds of cooperative behaviors needed to perpetuate niche-constructing activities and set the co-evolutionary relationships responsible for domestication into motion.

In addition to providing a more reliable supply of critical resources, however, resource management promotes notions of ownership of the resources and the manipulated environments in which they are grown, while also creating a real potential for uneven access to the rewards of collective activities [112]. These forces are evident in Near Eastern societies where, with agricultural emergence, we see a change from communal storage of vital resources in locations accessible to the whole community, to the development of household storage facilities within individual households outside the view of others, as well as the beginning of

increasing differentiation in access to basic resources [113,114]. The centrifugal tensions that arise with increased reliance on managed resources extracted from constructed niches make behaviors that combat them even more important to the survival of the group. These tensions, then, likely created additional pressures for behaviors that promote community cohesion and keep it from splintering into competing sub-groups.

Here is where NCT insights about the human capacity for language, symbolic communication, and information transfer through social learning can be seen to play a clear role in promoting the cooperative behaviors essential to the success of small-scale societies [115]. Rather than engaging in a continual process of calculation of self-interest and punishment of non-cooperators, humans, it is argued, have developed a means of maintaining cooperative behavior through systems of shared norms, customs, and values, that are played out and reinforced in an array of everyday and ritual activities, and that set unambiguous, widely acknowledge rules for cooperative behavior that make failure to abide by these rules expensive. These behaviors are on full display in societies on the threshold of plant and animal domestication which experience an uptick in social and ritual activities directed primarily at enhancing social cohesion and collective activities. In the Near East these activities are reflected in the construction of communal houses and megalithic structures, elaboration of symbolic lexicons, large scale feasting activities, and ritualized practice of day-to-day activities [116], all of which can be seen as vehicles for promoting the cooperative behaviors needed to perpetuate the collective activities and social cohesions that sustain these groups [117].

2.5 Rates of change

Niche-constructing activities are also held to have a profound impact on rates of evolutionary change, both speeding up and slowing down responses to selection [44,96]. Domestication provides a prime opportunity for exploring these impacts. Plastic responses to novel environments have a far greater potential for rapid evolutionary change than those induced through mutation. This is because mutation-induced novelties affect the survival of only one individual and must await a slow process of selection and transmission to spread more widely. In contrast, environmentally-induced novelties, such as those that arise in the context of niche

construction, affect numerous individuals, and can persist, even in the absence of strong positive selection, simply through the presence of the inducing factor [58,59]. Moreover, just as exposure to anthropogenic environments has been demonstrated to result in a dramatic increase in plastically-driven trait changes [61], anthropogenically-induced trait changes have also been shown to occur at a rate roughly twice as fast as changes stemming from non-human drivers. The rate of trait changes stemming from human harvest of wild populations is particularly rapid, estimated at about three times faster than other anthropogenically-induced trait changes [62]. Thus, the introduction of plants and animals to novel managed environments, whether they sought out these environments themselves or were relocated there by humans, is likely to have induced the expression of otherwise cryptic variation that would have resulted in rapid population-wide evolutionary change in receptive species. Periods of evolutionary stasis might be expected to follow initial introduction to anthropogenic environments as plants and animals living within these environments experience conditions of relaxed selection during which genetic variation accumulates in managed populations [57].

Humans, with their ability to spontaneously invent or alter goal-directed behaviors and to widely pass on effective behaviors, are particularly potent factors in setting the rate of evolutionary change [100]. These goal-directed behaviors can be expected to have had a profound impact on the rates of evolution in managed plants and animals, both as they respond to human management in ways that make them better equipped to benefit from their relationship with humans and as humans deliberately encourage traits of interest (even otherwise deleterious ones) and discourage those that make the partnership with the plant or animal less attractive. This is especially the case when humans begin to tinker with genomic reorganization by either encouraging the persistence of naturally occurring hybrids that have traits of interest, or becoming directly engaged in cross-breeding individuals or species for directed goals.

3. Contrasting neo-Darwinian and NCT informed explanatory frameworks

We can distill the fundamental differences between NCT informed perspectives and neo-Darwinian ones (briefly outlined in section 2 and explored at great length in 26 and 27) into two

contrasting explanatory frameworks for initial domestication that generate distinct predictions as to how the process unfolded (Table 2) [33]. A neo-Darwinian framework (Table 2: Column 3) begins with some event that causes an imbalance between human population levels and environmental carrying capacity [118,119]. Niche-constructing activities are only adopted as a way of compensating for the lower returns of less intensive resource extraction strategies and are considered a sign of a decrease in foraging efficiency [120]. Social and ritual activities, if mentioned at all, are portrayed as responses to tensions arising from resource depression and are directed toward increasing the competitive (and reproductive) advantage of some foragers over others [121,122]. And the co-evolutionary relationship between humans and target domesticates at the heart of the process of domestication unfolds at a gradual, incremental pace as the result of the accumulation of hundreds of decisions of individual foragers shaped by natural selection [22,25].

A NCT informed framework (Table 2: Column 4), in contrast, is set within environments in which abundant resources can be reliably found within circumscribed areas and population levels are below carrying capacity [28,33,111]. Niche-constructing activities should be evident from the outset and will likely intensify through time as they keep carrying capacity in balance with population [31]. Social and ritual activities play a key, catalytic role in NCT informed frameworks providing vehicles for transmitting environmental knowledge from one generation to the next, enhancing community cohesion, and combatting forces that encourage intra-group differences in access to basic resources [116]. Throughout, the process proceeds at an uneven pace, with periods of stasis punctuated by periods of rapid change, driven by both the rapid expression and accumulation of variation resulting from developmental processes and turbo-charged by the human capacity for inventing and transmitting goal-directed behaviors [34].

Not only can one draw clear, testable, predictions for these two contrasting frameworks, there is also a growing body of empirical data available to perform these tests (Table 2: Column 5), from enhanced climate proxies [123], ways of detecting human eco-system engineering [124], improved methods (morphometric and genetic) for tracing the process of domestication [125], large archaeobiological assemblages, and more precise direct dating techniques [126]. Domestication is, moreover, a global phenomenon that arose independently

in multiple areas around the world (Figure 1). Records for initial domestication in the Near East and eastern North America are particularly robust [111], those for China and the Andes are becoming increasingly well-resolved [28,127-129], and there are on-going efforts at documenting domestication in other world areas, including Mesoamerica, the Amazon, West and East Africa, and South East and Island Southeast Asia [130].

While systematic head-to-head evaluations of the different neo-Darwinian and NCT predictions about initial domestication are only just beginning to be performed [31,111], a review of empirical data from regions with the more complete records tend to support NCT generated predictions (see SI for a longer discussion). Mounting evidence from several world areas suggests that initial domestication took place in the context of small semi- to fully sedentary settlements strategically located at the junction of multiple resource-rich eco-zones in the absence of demonstrable resource pressure [28,31-33,111,127,131]. Niche constructing activities in each of these areas are evident well before any evidence of initial domestication [31,124,132]. Moreover, the uptick of social and ritual activities that accompanies increasing reliance on managed resources seems to be directed toward promoting community cohesion and combatting centrifugal tendencies toward competition [117,133]. And while the process of initial domestication and subsequent agricultural emergence unfolded over a millennium or more in most regions [32], the pace of change seems to be one in which long periods of relative stasis are punctuated by periods of more rapid change [26,34,126].

4. Domestication as a model system for the extended evolutionary synthesis

Beyond providing a tractable system for exploring the value of NCT as an explanatory framework for initial domestication, a case can also be made that domestication serves as an ideal model system for examining broader issues at the heart of the argument for extending the Modern Synthesis to accommodate recent developments in evolutionary biology, ecology, and the social sciences. Not only is NCT one of the cornerstones of the EES, as the above review of the relevance of NCT to initial domestication makes clear, essentially all the key concepts that fall under the EES umbrella are involved in the process. Recent advances in the study of initial domestication of plants and animals offer particularly promising potential avenues for empirical

evaluation of core EES concepts of reciprocal causation, constructive development, hierarchy in targets of selection, inclusive inheritance, and the tempo of evolutionary process highlighted in Table 1. Some of these avenues of future research are outlined here.

4.1 Causality

Adherents of neo-Darwinian informed explanatory frameworks tend to characterize domestication as a one-way adaptive response to new selective pressures [134] in which niche construction, and other developmental processes involved in domestication are portrayed either as adaptive responses to changing selective pressures [135], or as sources of environmental variability no different from stochastically derived variations in environmental background conditions that elicit adaptive responses [136]. This view stands in strong contrast to an EES view that envisions evolutionary events such as domestication as driven by “interacting bouts” of constructive development and selection that may actually lead and shape genetically encoded adaptive change rather than follow it [97]. The increasingly fine-grained record of initial domestication in multiple world areas provides an opportunity for assessing which of these contrasting views of the nature of causation and the primacy of natural selection in evolution is best supported by being able to more precisely track both selective and developmental processes involved in plant and animal domestication.

4.2 Directionality

All the constructive process that proponents of the EES maintain shape evolutionary trajectories in non-random, directional ways are in play in plant and animal domestication, with niche construction, developmental phenotypic plasticity, and genetic accommodation playing especially prominent roles [33]. Whole-genome sequencing has been accomplished for many of today’s domestic crops and livestock, as well as for many of their progenitors [137-140], providing a remarkably detailed understanding of the genetic basis for key domestication traits. Advances in the analysis of DNA extracted from archaeobiological remains and the ability to precisely date these remains, moreover, provides an excellent opportunity to monitor the interplay among these developmental processes and manifestations of heritable genetic

changes in emergent domesticates. Recent studies of ancient DNA extracted from ancient maize, for example, have made it possible to track the appearance of specific domestication genes in this important crop plant as it moved from the heartland of initial domestication in southwestern Mexico up through central Mexico and into the southwestern US [95,141,142]. Another recent study of ancient DNA extracted from horse skeletal remains finds evidence of enrichment for genes involved in androgen and steroid hormone receptor binding indicative of the selective pressures for behavioral and cognitive changes central to animal domestication [143]. This same study also found evidence for enrichment of genes affecting tissues and cell types derived from the neural crest that lend support for the hypothesis that the neural crest development is involved in the variety of associated traits commonly found in domestic animals [74].

New molecular techniques are coming on-line that make it possible to perform population-wide screenings for epigenetic markers [52]. Application of these techniques to directly dated archaeobiological remains might allow for an empirical assessment of the hypothesis advanced here that developmental plasticity plays a leading, perhaps dominant, role in shaping the genotype of domesticates.

4.3 Targets of selection

Domestication also provides a case study for exploring the targets of selection, especially the impact of selection operating above the level of individual alleles in driving evolutionary change. The possibility that the constellation of traits found in domestic animals arise through selection operating on neural crest cells [74] is one such opportunity for assessing the role of “exploratory behavior” among systems above the level of genes in facilitating evolutionary change [144]. The importance of the transmission of ecological inheritance and the reinforcement of cooperative behaviors in domestication, especially as they relate to human behavior, widens the focus of selection beyond gene frequencies of individuals to one that targets the distribution of traits across trans-generational populations, thus opening the way for an examination of the role of group selection in evolution [9,10,12,145].

4.4 Inheritance

As outlined above, domestication offers an opportunity to assess the impact of all the additional channels of information transfer that EES proponents maintain are equally as genetic inheritance in evolution, including epigenetic inheritance, ecological inheritance, social learning and culture. Consideration of the domestication process promises new insights into the role of these inclusive or “soft” forms of inheritance in shaping evolutionary change [9,10,146].

The importance of culture as a form of soft inheritance is particularly noteworthy. Evolutionary biologists operating within the framework of the MS have traditionally had a very difficult time accounting for the role of culture in evolutionary change [145,147]. Human culture is often relegated to the status of a proximate causal process that might help “fine-tune” the variation on which selection (the ultimate cause of evolutionary change) acts, but that has little influence on directional change in evolution [148,149]. This view is embraced by archaeologists working within MS informed frameworks who dismiss the role of human agency or intentionality in evolutionary change – biological or cultural [20,22,24]. This discomfort might also be a reason that many social scientists and scholars working within the humanities tend to down-play or even dismiss the role of human/environmental interaction, and the relevance of evolutionary biology, focusing instead on more purely social, ritual, and symbolic behaviors in explanatory frameworks of domestication and culture change in general [7,33]. A view of culture as a form of soft inheritance builds a bridge between the biological and social sciences, and the humanities, that neither negates or downplays the role of culture in evolution, nor glorifies it as something exceptional that operates outside of nature. Recognizing culture and other human cognitive abilities as extensions of the use of acquired knowledge and social learning by non-human animals to enhance the fidelity of information transmitted along external inheritance channels, brings these behaviors squarely within the parameters of the evolutionary process, while also acknowledging the unique and powerful parts they play in shaping the evolutionary trajectory of humans and biota affected by human actions.

The niche construction literature has made significant contributions to documenting the role of domestication in shaping gene-culture interactions and their effect on human biological evolution, with the study of the development of lactose tolerance and resistance to sickle cell

anemia in humans practicing dairying and yam cultivation primary examples of this important work [150]. In addition to having had some impact on the human genome, however, domestication has also played a major role in influencing the course of cultural change, thus providing an opportunity for exploring the importance of human goal-directed behavior and cultural transmission in human cultural evolution. Consideration of domestication as a model system for evaluating core concepts of the EES also allows for an examination of the role of human agency and intentionality in shaping the evolutionary trajectories of partner species in domesticatory relationships, as well as in the evolution of other non-participant organisms affected by these behaviors. Moreover, a view that sees target species as assuming an active part in directing the course of domesticatory relationships provides an opportunity for examining the reciprocal impact they have in shaping the trajectory of human cultural and biological evolution.

5.5 Tempo and Pace of Change

Finally, the ability to set the process of domestication in an increasingly well-constrained temporal framework is perhaps one of the most appealing aspects of domestication as a model system for exploring core assumptions of the extended synthesis. Advocates of the EES sometimes lament the lack of empirical examples of the role of genetic accommodation in fixing traits arising from developmental processes, maintaining that the signs of this process may disappear so quickly from a natural population as to make it appear as if these traits arose through standard selection operating on genetic variants following the MS paradigm [52]. Possible systems suggested for documenting these fast-moving processes include natural environments undergoing dramatic rapid modification through climate change or anthropogenically-induced habitat destruction [85], invasive species moving into and adapting to novel environments [52,53], and comparisons of the capacities for the expression of phenotypic plasticity in extant ancestral populations with those of derived ones [57].

Domestication resembles these potential model systems in that it involves the introduction and subsequent adaptation of species to novel anthropogenically modified environments that provide optimal settings for assessing the role of constructive processes in

the genetic fixation of adaptive traits. In many cases, there are living representatives of both domesticates and their progenitors that might allow for an assessment of the extent to which ancestral plasticity served as a source for fixed genetic traits in their domestic descendants, and thus to assess whether these traits arose as a response to selective pressures, or through the plastic expression of existing previously cryptic traits that were subsequently fixed through the process of genetic accommodation [57]. Perhaps even more exciting is the potential of ancient DNA for determining the timing of the sequence of plastic expression and genetic accommodation behind the fixation of domestication traits in “real time” as it unfolded during the process of initial domestication. Domestication is then a model system that combines multiple interacting constructive processes and multiple internal and external inheritance systems resulting in significant evolutionary change in multiple species, set within an increasingly well-resolved temporal sequence, that seems to meet the requirements for a gold standard system EES advocates believe is needed for a rigorous evaluation of need for revision of standard evolutionary theory [60].

6. Conclusions

Darwin used domestication as a model system for exploring mechanisms of variation in plants and animals under human control to support his thesis of evolution through natural selection [151]. Experimental cross breeding of domesticates (both plants and animals) also provided model systems that allowed Greger Mendel to discover the rules of heredity responsible for the transmission of variation across generations that provided the fuel for Darwinian selection. Domesticates and the process of domestication were then foundational systems for the modern evolutionary synthesis that brought Darwinian theory of evolution together with Mendelian genetics through the application of population genetics [1-3]. It seems only appropriate that those seeking to build on this important foundation turn once again to domestication as a model system for assessing how recent insights into the role of multiple shaping processes and forms of inheritance should be incorporated into and extended understanding of evolution.

References

- 1 Huxley J. 1942 *Evolution: The Modern Synthesis*. New York: Harper and Brothers.
- 2 Mayr E. 1942 *Systematics and the origin of the species from the viewpoint of a zoologist*. New York NY: Columbia University Press.
- 3 Simpson GG. 1977 *Tempo and mode in evolution*. New York: Columbia University Press.
- 4 Gould S J, Lewontin RC. 1979 The spandrels of San Marco and the Panglossian paradigm: A critique of the adaptationalist programme. *Procs Ry Soc Lon B* 205, 581–598.
- 5 Eldredge N, Gould SJ. 1972 Punctuated equilibria: An alternative to phyletic gradualism. In: *Models in Paleobiology* (ed TJ Schopf), pp. 82–115. San Francisco, CA: Freeman, Cooper.
- 6 Gould SJ. 2002 *The Structure of Evolutionary Theory*, Cambridge, MA: Belknap.
- 7 Odling-Smee FJ, Laland KN, Feldman MW. 2003. *Niche construction: the neglected process in evolution*. Monographs in population biology 37, Princeton, NJ: Princeton University Press.
- 8 West-Eberhard MJ. 2003 *Developmental plasticity and evolution*. Oxford, UK: Oxford University Press.
- 9 Jablonka E, Lamb MJ. 2010 Transgenerational Epigenetic inheritance. In *Evolution: The Extended Synthesis* (eds M Pigliucci, GD Müller), pp. 137-174. Cambridge, MA: MIT Press.
- 10 Jablonka E, Lamb MJ. 2008 Soft inheritance: challenging the modern synthesis. *Genet. Mol. Biol* 31, 389–395.

- 11 Pigliucci M, Müller G. (eds) 2010 *Evolution: the extended synthesis*. Cambridge, MA: The MIT Press.
- 12 Laland KM, Uller T, Feldman MW, Sterenly K, Müller G, Moczek A. Jablonka E, Odling-Smee J. 2015 The extended evolutionary synthesis: its structure, assumptions and predictions. *Proc. R. Soc. B Lond* 282, 20151019.
- 13 Scott-Phillips TC, Laland KN, Shuker DM, Dickins TE, West SA. 2014 The niche-construction perspective. A critical appraisal. *Evolution* 68, 1231–1243.
- 14 Futuyma D. 2017 The extended synthesis today: extend or amend? *Interface Focus* [citation is from his oral presentation. not certain whether DF is contributing to the volume and if so what title of his contribution will be.]
- 15 Lande R. 2017. Evolution of phenotypic plasticity. *Interface Focus* [citation is from his oral presentation. not certain whether RL is contributing to the volume and if so what title of his contribution will be.]
- 16 Schlichting CD, Wund MA. 2014 Phenotypic plasticity and epigenetic marking: an assessment of evidence for genetic accommodation. *Evolution* 68, 656-672.
- 17 Smith BD, Zeder MA. 2013 The onset of the Anthropocene. *Anthropocene* 4, 8–13.
- 18 Dunnell R. 1980 Evolutionary theory and archaeology. *Advances in Archaeological Method and Theory* 3, 35–99.
- 19 O’Brien MJ, Lyman RL. 2000 *Applying Evolutionary Archaeology*. New York: Kluwer Academic/Plenum.

- 20 Lyman RL, O'Brien MJ. 1998 The goals of evolutionary archaeology: history and explanation. *Curr Anthropol* 39, 615–652.
- 21 Rindos D 1985 Darwinian selection, symbolic variation, and the evolution of culture. *Curr Anthropol* 26, 65–77.
- 22 Rindos D. 1984 *The origins of agriculture: an evolutionary perspective*. Orlando, FL: Academic Press.
- 23 Hawkes K, O'Connell JF. 1992 On optimal foraging models and subsistence transitions. *Curr Anthropol* 33, 63–65.
- 24 Gremillion KJ, Barton L, Piperno D. 2014 Particularism and the retreat from theory in the archaeology of agricultural origins. *Proc Natl Acad Sci USA* 111, 6171–6177.
- 25 Winterhalder B, Kennett DJ (2006) Behavioral ecology and the transition from hunting and gathering to agriculture. In *Behavioral ecology and the transition to agriculture* (eds DJ Kennett, B Winterhalder), pp. 1–21. Berkeley, CA: University of California Press.
- 26 Zeder MA. 2009 Evolutionary biology and the emergence of agriculture: the value of co-opted models of evolution in the study of culture change. In: *Macroevolution in human prehistory* (eds A Prentiss, I Kuijt, J Chatters), pp. 157–210. New York, NY: Springer.
- 27 Zeder MA. 2012 The Broad Spectrum Revolution at 40: resource diversity, intensification, and an alternative to optimal foraging explanations. *J Anthropol Archaeol* 31, 241–264.

- 28 Crawford G. 2011 Advances in understanding early agriculture in Japan. *Curr Anthropol* 52, S331– S345.
- 29 Smith BD. 2011 General patterns of niche construction and the management of wild plant and animal resources by small-scale pre-industrial societies. *Philos Trans R Soc Lond B Biol Sci* 366, 836–848.
- 30 Smith BD. 2012 A cultural niche construction theory of initial domestication. *Biol Theory* 6, 260–271.
- 31 Smith BD. 2016 Neo-Darwinism, niche construction theory, and the initial domestication of plants and animals. *Evol Ecol* 30, 307-324.
- 32 Zeder MA. 2015 Core concepts in domestication research. *Proc Natl Acad Sci USA* 112, 3191–3198.
- 33 Zeder MA. 2016 Domestication as a model system for niche construction theory. *Evol Ecol* 30,325-348.
- 34 Kuijt I, Prentiss AM. 2009 Niche construction, macroevolution, and the Late Epipaleolithic of the Near East. In *Macroevolution in human prehistory* (eds AM Prentiss, I Kuijt, J Chatters), pp. 253–271. New York, NY: Springer.
- 35 Laland KN, Odling-Smee FJ, Feldman MW, Kendall JR. 2009 Conceptual barriers to progress within evolutionary biology. *Found Sci* 4, 195–216.
- 36 Laland K, Matthews B, Feldman MW. 2016 An introduction to niche construction theory. *Evol Ecol* 30, 191-202.

- 37 Antón SC, Potts R, Aiello LC (2014) Evolution of early Homo: an integrated biological perspective. *Science* 345(6192), 1236828.
- 38 Fuentes A, Wyczalkowski M, MacKinnon KC. 2010 Niche Construction through Cooperation: A Nonlinear Dynamics Contribution to Modeling Facets of the Evolutionary History in the Genus Homo. *Curr Anthropol* 51(3), 435-444.
- 39 Kendal J, Tehrani JJ, Odling-Smee FJ. 2011 Human niche construction in interdisciplinary focus. *Philos Trans R Soc B* 366, 785–792.
- 40 Flynn E, Laland KN, Kendal R et al (2013) Developmental niche construction. *Dev Sci* 16, 296–313.
- 41 Post DM, Palkovacs EP. 2009 Eco-evolutionary feedbacks in community and ecosystem ecology: interactions between the ecological theatre and the evolutionary play. *Philos Trans R Soc B* 364, 1629–1640.
- 42 Hendry AP. 2017 *Eco-evolutionary dynamics*. Oxfordshire, UK: Princeton University Press.
- 43 Laland KN, Boogert NJ. 2008 Niche construction, co-evolution, and biodiversity. *Ecol Econ* 69, 731–736.
- 44 Odling-Smee FJ, Erwin D, Palkovacs E, Feldman M, Laland KN. 2013 Niche construction theory: a practical guide for ecologists. *Q Rev Biol* 88, 3–28.
- 45 Matthews B, de Meester L, Jones CG, Ibelings BW, Bouma TJ, Nuutinen V, de Koppel J, Odling-Smee J. 2014 Under niche construction: an operational bridge between ecology, evolution, and ecosystem science. *Ecol Monogr* 84(2), 245–263.
- 46 Zeder MA. 2012 The domestication of animals. *J Anthropol Res* 68, 161–190.

- 47 Smith BD. 2006 Documenting plant domestication in the archaeological record. In *Documenting domestication: new genetic and archaeological paradigms* (eds MA Zeder, E Emshwiller, BD Smith, D Bradley), pp. 15–24. Berkeley, CA: University of California Press.
- 48 Duarte CM, Marbá N, Holmer H. 2007 Rapid domestication of marine species. *Science* 316, 382–83.
- 49 Donohue K. 2014 Why ontogeny matters during adaption: developmental niche construction and pleiotropy across the life cycle in *Arabidopsis thaliana*. *Evolution* 68, 32–47.
- 50 Fuller DQ, Allaby R. 2009 Seed dispersal and crop domestication: shattering, germination and seasonality in evolution under cultivation. *Annu Plant Rev* 38, 238–295.
- 51 Larson G, Fuller DQ. 2014 The evolution of animal domestication. *Annu Rev Ecol Evol Syst* 45, 115–136.
- 52 Pigliucci M. 2010. Phenotypic plasticity. In *Evolution: The Extended Synthesis*. (eds M Pigliucci, GD Müller), pp. 354-378. Cambridge, MA: MIT Press.
- 53 Snell-Rood EC. 2012 Selective processes in development: implications for the costs and benefits of phenotypic plasticity. *Integr Comp Biol* 52, 31–42.
- 54 Gremillion KJ, Piperno DR. 2009 Human Behavioral Ecology, phenotypic (developmental) plasticity, and agricultural origins: insights from the emerging evolutionary synthesis. *Curr Anthropol* 50, 615–619.

- 55 Piperno DR, Holst I, Winter K, McMillan O. 2015 Teosinte before domestication: experimental study of growth and phenotypic variability in Late Pleistocene and early Holocene environments. *Quat Int* 363,65-77.
- 56 Asouti E. 2017. Human palaeoecology in southwest Asia during the early pre- pottery Neolithic (9700-8500 cal BC): the plant story. In: *The Construction of Neolithic Corporate Identities*. (eds M Benz, HG Gebel, T Watkins), pp. in press. Proceedings of the 9th ICAANE Workshop, SENSPSE 18. Berlin: ex oriente.
- 57 Moczek AP, Sultan S, Foster S, Ledón-Rettig C, Dworkin I, Nijhout HF, Abouheif E, Pfennig D. 2011 The role of developmental plasticity in evolutionary innovation. *Proc Biol Sci Lond B* 278, 2705–2713.
- 58 West-Eberhard MJ. 2005 Developmental plasticity and the origin of species differences. *Proc Natl Acad Sci USA* 102, 6543–6549.
- 59 West-Eberhard MJ. 2005 Phenotypic accommodation: adaptive innovation due to developmental plasticity. *J Exp Zool B Mol Dev Evol B* 304, 610–618.
- 60 Snell-Rood EC, Van Dyken JD, Cruickshank TE, Wade MJ, Moczek AP. 2010 Toward a population genetic framework of developmental evolution: costs, limits, and consequences of phenotypic plasticity. *Bioessays* 32, 71–81.
- 61 Hendry AP, Farrugia TJ, Kinnison MT. 2008 Human influences on rates of phenotypic change in wild animal populations. *Mol Ecol* 17, 20–29.
- 62 Palkovacs EP, Kinnison MT, Correa C, Dalton CM, Hendry AP. 2011 Fates beyond traits ecological consequences of human induced trait change. *Evol Appl* 5, 183-191.
- 63 Darimont CT, Carlson SM, Kinnison MT, Paquette PC, Reimchena TE, Wilmers CC. 2009

- Human predators outpace other agents of trait change in the wild. *Proc Natl Acad Sci USA* 106(3), 952-954.
- 64 Coltman DW, O'Donoghue P, Jorgenson JY, Hogg JT, Strobeck C, Festa-Blanchet M. 2003 Undesirable evolutionary consequences of trophy hunting. *Nature* 426, 655-658.
- 65 Gremillion KJ. 1993 The evolution of seed morphology in domesticated *Chenopodium*: an archaeological case study. *J Ethnobiol* 13, 149–169.
- 66 Wilcox G, Fornite S, Herveux L. 2008 Early Holocene cultivation before domestication in northern Syria. *Veg Hist Archaeobot* 17, 313–325.
- 67 Donohue K 2014 The epigenetics of adaptation: focusing on epigenetic stability as an evolving trait. *Evolution* 68, 617-619.
- 68 Allaby 2014 Domestication syndrome in plants. In *Encyclopedia of global archaeology*, (ed C Smith), pp. 2182-2184. New York, NY: Springer.
- 69 Jensen P. 2006 Domestication—from behavior to genes and back again. *Appl Anim Behav Sci* 97, 3–15.
- 70 Poncet V, Martel E, Allouis S, Devos K, Lamy F, Sarr A, Robert T. 2002 Comparative analysis of QTLs affecting domestication traits between two domesticated × wild pearl millet (*Pennisetum glaucum* L., *Poaceae*) crosses. *Theor Appl Genet* 104, 965-975.
- 71 Koinange EMK, Singh SP, Gepts P. 1996 Genetic control of the domestication syndrome in common bean. *Crop Sci* 36, 1037-1045.
- 72 Albert FW, Somel M, Carneiro M, Aximu-Petri A, Halbwax M, Thalmann O, Blanco-Aguilar JA, Plyusnina IZ, Trut L, Villafuerte R, et al. 2012 A comparison of brain expression levels in domesticated and wild animals. *PLoS Genet* 8, 1–16.

- 73 Carneiro M, Rubin C-J, Di Palma F, Albert FW, Alföldi J, Martinez Barrio A, Pielberg G, Rafati N, Sayyab S, Turner-Maier J, et al. 2014 Rabbit genome analysis reveals a polygenic basis for phenotypic change during domestication. *Science* 345, 1074–1079.
- 74 Wilkens AS, Wrangham RW, Tecumseh Fitch W. 2014 The “Domestication Syndrome” in mammals: a unified explanation based on neural crest cell behavior and genetics. *Genetics* 197, 795–808.
- 75 Conrad JL, Weinersmith KL, Brodin T, Saltz JB, Sih A. 2011 Behavioural syndromes in fishes: a review with implications for ecology and fisheries management. *J Fish Biol* 78, 395-435.
- 76 Stebbins JGL. 1951 Cataclysmic Evolution. *Scientific American*. 184, 54–59.
- 77 Shapiro JA. 2017 Biological action in read-write genome evolution. *J R Soc Interface* in press. [citation is from his oral presentation. not certain whether DF is contributing to the volume and if so what title of his contribution will be.]
- 78 Emschwiller E. 2006 Origins of polyploid crops: the example of the octoploid tuber crop *Oxalis tuberosa*. In: *Documenting domestication: new genetic and archaeological paradigms* (eds MA Zeder, DG Bradley, E Emschwiller, BD Smith), pp. 153-168. Berkeley, CA: University of California. Press.
- 79 Eriksson J, Larson G, Gunnarsson U, Bed’hom B, Tixier-Boichard M, Strömstedt L, Wright D, Jungerius A, Vereijken A, Randi E, Jensen P, Andersson L. 2008 Identification of *yellow skin gene* reveals a hybrid origin of the domestic chicken. *PLoS Genet* 4, e1000010.

- 80 Wheeler JC, Chickhi L, Bruford MW. 2006 Genetic analysis of the origins of South American camelids. In *Documenting Domestication: New Genetic and Archaeological Paradigms* (eds MA Zeder, DG Bradley, E Emshwiller, BD Smith), pp. 329–341. Berkeley, CA: University of California Press.
- 81 Weiss E, Kislev M, Hartmann A. 2006 Autonomous cultivation before domestication. *Science* 312, 1608–1610.
- 82 Tanno K-I, Wilcox G. 2006. How fast was wild wheat domesticated? *Science* 311, 1886.
- 83 Faris JD. 2014 Wheat domestication: key to agricultural revolutions past and present. In *Genomics of plant genetic resources* (eds R Tuberosa, A Graner, E Frison), pp. 439–464. Netherlands: Springer.
- 84 Pfennig DW, Wund MA, Snell-Rood EC, Cruickshank T, Schlichting CD, Moczek AP. 2010 Phenotypic plasticity's impacts on diversification and speciation. *Trends Ecol Evol* 25, 459–467.
- 85 Moczek AP. 2012 The nature of nurture and the future of evodevo: toward a comprehensive theory of developmental evolution. *Integrative Comp Biol* 52, 108–119.
- 86 Hillman, GC, Davies MS. 1992 Domestication rates in wild wheats and barley under primitive cultivation: preliminary results and archaeological implications of field measurements of selection coefficient. In *Préhistoire de l'agriculture: nouvelles approches expérimentales et ethnographiques*. (ed PC Anderson), pp. 113–158. Paris: Centre National de la Recherche Scientifique.

- 87 Maeda O, Lucas L, Silva F, Tanno K-I, Fuller DQ. 2016 Narrowing the harvest: increasing sickle investment and the rise of domesticated cereal agriculture in the Fertile Crescent. *Quat Sci Rev* 145, 226-237.
- 88 Zeuner FE. 1955. *A history of domesticated animals*. London: Hutchinson.
- 89 Lande R. 2009 Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *J Evol Biol* 22, 1435–46.
- 90 Fuller D, Lucas L. 2017 Adapting crops, landscapes, and food choices: Patterns in the dispersal of domesticated plants across Eurasia. In *From colonization to globalization: species movement in human history* (eds N. Boivin, R Crassard, M Petraglia), pp. in press. UK: Cambridge University Press.
- 91 Jones H, Leigh FJ, Mackay I, Bower MA, Smith LMJ, Charles MP, Jones G, Jones MK, Brown TA, Powell W. 2008 Population-based resequencing reveals that the flowering time adaptation of cultivated barley originated east of the fertile crescent. *Mol Biol Evol* 25, 2211-2219.
- 92 d'Alpoim Guedes J, Lu H, Li Y, Spengler RN, Wu X, Aldenderfer MS. 2013 Moving agriculture onto the Tibetan Plateau: the archaeobotanical evidence. *Archaeol Anthropol Sci* 6, 255-269.
- 93 Smith BD. 2017 Tracing the initial diffusion of maize in North America In *From colonization to globalization: species movement in human history* (eds N. Boivin, R Crassard, M Petraglia), pp. in press. UK: Cambridge University.
- 94 da Fonseca RR, Smith BD, Wales N, Cappellini E, Skoglund P, Fumagalli M, Samaniego JA, Carøe C, Ávila-Arcos MC, Hufnagel DE, Korneliussen TS, Vieira FG, Jakobsson M, Arriaza B, Willerslev E, Nielsen R, Hufford MB, Albrechtsen A, Ross-Ibarra J, Gilbert MT. 2015

- The Origin and Evolution of Maize in the Southwestern United States. *Nature Plants* 1(1).
- 95 Jaenicke-Després V, Buckler ES, Smith BD, Gilbert MTP, Cooper A, Doebley J, Pääbo. 2003 Early allelic selection in maize as revealed by ancient DNA. *Science* 302, 1206–1208.
- 96 Erwin DH. 2008 Macroevolution of ecosystem engineering, niche construction and diversity. *Trends Ecol Evol* 23, 304–310.
- 97 Laland KN, Odling-Smee FJ, Gilbert SF. 2008 Evo-Devo and niche construction: building bridges. *J Exp Zool Part B Mol Dev Evol* 310B, 549–566.
- 98 Odling-Smee FJ. 2010 Niche inheritance. In *Evolution: the extended synthesis* (eds M Pigliucci, GB Müller), pp. 175–207. Cambridge, MA: MIT Press.
- 99 Odling-Smee FJ, Laland KN. 2012 Ecological inheritance and cultural inheritance: what are they and how do they differ? *Biol Theory* 6(3), 220-230.
- 100 Laland KN, Sterelny K. 2006 Seven reasons (not) to neglect niche construction. *Evolution* 60, 1751–1762.
- 101 Dean LG, Kendal RL, Schapiro SJ, Thierry B, Laland KN. 2013 Identification of the social and cognitive processes underlying human cumulative culture. *Science* 335, 1114–1118.
- 102 Berkes F. 2008 *Sacred ecology*, 2nd edn. New York, NY: Routledge.
- 103 Anderson MK. 2005 *Tending the wild*. Berkeley, CA: University of California Press.
- 104 Deur D, Turner N. (eds) 2005 *Keeping it living: traditions of plant use and cultivation on*

- the Northwest Coast of North America*. Seattle, WA: University of Washington Press.
- 105 Hunn E. 1993 What is traditional ecological knowledge? In: *Traditional ecological knowledge: wisdom for sustainable development* (eds NM Williams, G Baines), pp. 13–15. Canberra, AU: Centre for Resource and Environmental Studies, Australian National University.
- 106 Bliege-Bird R, Bird DW, Copping BF, Parker CH, Jones JH. 2008 The “fire stick farming” hypothesis: Australian Aboriginal foraging strategies, biodiversity, and anthropogenic fire mosaics. *Proc Nat Acad Sci USA* 105, 14796–14801.
- 107 Krakauer DC, Page KM, Erwin D. 2009 Diversity, dilemmas, and monopolies of niche construction. *Am Nat* 173, 26–40.
- 108 Bettinger RL, Barton L, Morgan C. 2010 The origins of food production in north China: a different kind of agricultural revolution. *Evol Anthropol* 19, 9–21.
- 109 Lehmann L. 2008 The adaptive dynamics of niche-constructing traits in spatially subdivided populations: evolving posthumous extended phenotypes. *Evolution* 62, 549–566.
- 110 Lehmann L. 2007 The evolution of trans-generational altruism: kin selection meets niche construction. *J Evol Biol* 20, 181–189.
- 111 Zeder MA, Smith BD. 2009 A conversation on agriculture: talking past each other in a crowded room. *Curr Anthropol* 50, 681–691.
- 112 Bowles S, Choi J-K. 2013 Co-evolution of farming and private property during the early Holocene. *Proc Natl Acad Sci USA* 110, 8830–8835.

- 113 Kuijt I 2000 People and space in early agricultural villages: Exploring daily lives, community size, and architecture in the late Pre-Pottery Neolithic. *J Anthropol Archaeol* 19 75–102.
- 114 Kuijt I, Finlayson B. 2009 Evidence for food storage and pre-domestication granaries 11,000 years ago in the Jordan Valley. *Proc Natl Acad Sci USA* 106, 10966-10970.
- 115 Sterelny K. 2007 Social intelligence, human intelligence and niche construction. *Philos Trans R Soc B* 362, 719–730.
- 116 Zeder MA. 2012 Religion and the revolution: the legacy of Jacques Cauvin. *Paléorient* 37(1), 39–60.
- 117 Kuijt I. ed 2000 *Life in Neolithic farming communities: social organization, identity, and differentiation*. New York, NY: Plenum.
- 118 Piperno DR. 2006 The origins of plant cultivation and domestication in the Neotropics: a behavioral ecological approach. In: *Behavioral ecology and the transition to agriculture* (eds DJ Kennett, B Winterhalder), pp. 137–166. Berkeley, CA: University of California Press.
- 119 Stutz AJ, Munro ND, Bar-Oz G. 2009 On increasing the resolution of the Broad Spectrum Revolution in the southern Levantine Epipaleolithic (19–12 ka). *J Hum Evol* 56, 294–306.
- 120 Munro ND, Atici L. 2009 Human subsistence change in the Late Pleistocene Mediterranean Basin: the status of research on faunal intensification, diversification, and specialization. *Before Farming: The Archaeology of Old World Hunter–Gatherers* 2009/1, article 1.

- 121 Aldenderfer, M., 2006. Costly signaling, the sexual division of labor, and animal domestication in the Andean Highlands. In *Behavioral Ecology and the Transition to Agriculture*. (eds DJ Kennett, B Winterhalder), pp. 167–196. Berkeley, CA: University of California Press.
- 122 Starkovich BM., Stiner MC. 2009 Hallan Çemi Tepesi: high ranked game exploitation alongside intensive seed processing at the Epipaleolithic Neolithic transition in southeastern Turkey. *Anthropozoologica* 44, 41–61.
- 123 Roberts N, Eastwood WJ, Kuzucuoglu C, Fiorentino G, Carcuta V. 2011 Climatic, vegetation and cultural change in the eastern Mediterranean during the mid-Holocene environmental transition. *Holocene* 21, 147–162.
- 124 Asouti E, Kabukcu C, White CE. 2015 Early Holocene woodland vegetation and human impacts in the arid zone of the southern Levant. *Holocene*. 25, 1565-1580.
- 125 Zeder MA, Emshwiller E, Smith BD, Bradley DG. 2006 Documenting domestication: the intersection of genetics and archaeology. *TIG* 22, 139-55.
- 126 Byrd B. 2005. Reassessing the emergence of village life in the Near East. *J Archaeol Res* 13, 231–290.
- 127 Ren X, Lemoine X, Mo D, Kiffer TR, Guo Y, Qin Z, Liu X. 2016 Foothills and intermountain basins: does China’s fertile arch have ‘Hilly Flanks? *Quat Int* 334+335, 165-178.
- 128 Mengoni-Goñalons G, Yacobaccio H. 2006 The domestication of South American camelids: A view from the South-Central Andes. In *Documenting Domestication: New Genetic and Archaeological Paradigms* (eds MA Zeder, DG Bradley, E Emshwiller, BD Smith), pp. 228–244. Berkeley, CA: University of California Press.

- 129 Pearsall D. 2008 Plant domestication and the shift to agriculture in the Andes. In *Handbook of South American Archaeology* (eds H Silverman, WH Isbell), pp. 105-120. New York, NY:Springer.
- 130 Smith BD. 1998 *The emergence of agriculture*. New York, NY: Scientific American Library.
- 131 Smith BD. 2009 Resource Resilience, Human Niche Construction, and the Long-Term Sustainability of Pre-Columbian Subsistence Economies in the Mississippi River Valley Corridor. *J Ethnobiol* 29(2), 167-183.
- 132 Aceituno FJ, Loaiza N, Delgado-Burbano M, Barrientos G. 2013 The initial settlement of Northwest South America during the Pleistocene–Holocene transition: synthesis and perspectives. *Quatern Int* 30, 23–33.
- 133 Zeder MA, Spitzer MD. 2016. New insights into broad spectrum communities of the Early Holocene Near East: the birds of Hallan Çemi. *Quat Sci Rev* 151, 140-159.
- 134 Williams GC. 1992 Gaia, nature worship, and biocentric fallacies. *Q Rev Biol* 67, 479–486.
- 135 Coding B, Bird D. 2015 Behavioral ecology and the future of archaeological science. *J Archaeol Sci* 56, 9-20.
- 136 Broughton JM, Cannon MD, Barelink EJ. 2010 Evolutionary ecology, resource depression, and niche construction theory: applications to central California hunter-gatherers and Mimbres-Mogollon agriculturalists. *J Archaeol Method Theory* 17, 317–421.
- 137 Benchley R, Spannagl M, Pfeifer M, Barker GLA, D’Amore R, Allen AM, McKenzie N, Kramer M, Kerhornou A, Bolser D, Kay S, Waite D, Trick M, Bancroft I, Gu Y, Huo N, Luo

- M-C, Sehgal S, Kianian S, Gill B, Anderson O, Kersey P, Dvork J, McCombie R, Hall A, Mayer KFX, Edwards KJ, Bevan MW, Hall N. 2012. Analysis of the bread wheat genome using whole genome shotgun sequencing *Nature* 491,705-710.
- 138 Groenen MAM. 2016 A decade of pig genome sequencing: a window on pig domestication and evolution *Genet Sel Evol* 48, 23.
- 139 Rubin C-J, Zody MC, Eriksson J, Meadows JRS, Sherwood³ E, Webster MT, Jiang L, Ingman M, Sharpe T, Ka S, Hallböök F, Besnier F, Carlborg Ö, Bed'hom B, Tixier-Boichard M, Jensen P, Siegel P, Lindblad-Toh K, Andersson L. 2010 Whole-genome resequencing reveals loci under selection during chicken domestication *Nature* 464, 587-591.
- 140 Weber AL, Briggs WH, Rucker J, Baltazar BM, de Jesús Sánchez-Gonzalez J, Feng P, Buckler ES, Doebley J 2008. The genetic architecture of complex traits in teosinte (*Zea mays ssp. parviglumis*): new evidence from association mapping. *Genetics* 180, 1221–1232.
- 141 Ramos-Madriral J, Smith BD, Moreno-Mayar JV, Gopalakrishnan S, Ross-Ibarra J, Gilbert MTP, Wales N. 2016 Genome sequence of a 5,310-year-old maize cob provides insights into the early stages of maize domestication. *Curr Biol* 26(23), 3195-3201.
- 142 Vallebuena-Estrada M, Rodríguez-Arévalo I, Rougon-Cardoso A, Martínez González J, García Cook A, Montiel R, Vielle-Calzada J-P. 2016 The earliest maize from San Marcos Tehuacán is a partial domesticate with genomic evidence of inbreeding. *Proc Nat Acad Sci USA* 113, 14151-14156.
- 143 Librado P, Gamba C, Gaunitz C, de Sarkissian C, Pruvost M, Albrechtsen A, Fages A, Khan N, Schubert M, Jagannathan V et al. 2017. Ancient genomic changes associated with the domestication of the horse. *Science* 356, 442-445.

- 144 Gerhart JC, Kirschner MW. 2007 The theory of facilitated variation. *Proc Natl Acad Sci USA* 104, 8582 – 8589.
- 145 Wilson DS. 2010 Multilevel Selection and major transitions. In *Evolution: The Extended Synthesis* (eds M Pigliucci, GD Müller), pp. 81-93. Cambridge, MA: MIT Press.
- 146 Dickins T, Rahman Q. 2012 The extended evolutionary synthesis and the role of soft inheritance in evolution. *Proc. R. Soc. Lond. B* 278, 1721 – 1727.
- 147 Laland KN 2017 *Darwin's Unfinished Symphony: How culture made the human mind*. Princeton NJ; Princeton University Press.
- 148 Laland KN, Odling-Smee J, Hoppitt W, Uller T. 2013. More on how and why: cause and effect in biology revisited. *Biol Philos*. DOI 10.1007/s10539-012-9335-1
- 149 Laland KN, Uller T, Feldman MW, Sterelny K, Müller GB, Moczek A, Jablonka E, Odling-Smee FJ. 2014a Does evolutionary theory need a rethink? Yes, urgently. *Nature* 514, 161–164.
- 150 O'Brien MJ, Laland K. 2012 Genes, culture, and agriculture: an example of human niche construction. *Curr Anthropol* 53, 434–470.
- 151 Darwin C. 1868 *The variation of animals and plants under domestication*. UK: John Murphey.

Table Captions

Table 1: Core Assumptions of the Modern Synthesis and the Extended Evolutionary Synthesis – after Laland et al. [1]

Table 2: Test Implications and data sets of explanatory frameworks for initial domestication

Figure Caption

Figure 1: Currently recognized areas of domestication of plants and animals with some examples of plants and animals domesticated in each center. (1) Eastern North America: chenopodia, squash, sunflower, knotweed, and maygrass; (2) Southwest US: turkeys (3) Mesoamerica: maize, squash, beans, and turkeys; (4) northern Peru/Ecuador: squash and lima beans; (5) Amazonia: manioc, yams, peanuts, and Muscovy duck; (6) Andes: oca, potato, quinoa, amaranth, llama, alpaca, and guinea pigs; (7) sub-Saharan Africa: pearl millet, sorghum, and African rice; (8) Horn of Africa/Nile Valley: asses, tef; (9) Near East: wheat, barley, lentils, peas, sheep, goats, taurine cattle, and pigs; (10) Central Asia: horses, golden hamster; (11) South Asia: browntop millet, water buffalo, and zebu cattle; (12) North China: foxtail and broomcorn millet; (13) South China/Southeast Asia: rice and chickens; (14) Japan: barnyard millet, mung bean, burdock; and (15) New Guinea: bananas, yams, and taro. Note: list of domesticates not exhaustive.


Table 1: Core Assumptions of the Modern Synthesis and the Extended Evolutionary Synthesis – after Laland et al. [12]

	Modern Synthesis	Extended Evolutionary Synthesis
Causality	Natural selection is the pre-eminent cause of evolution, responsible for sorting variation on basis of adaptive fitness; operates as asymmetrical, unidirectional process in which organisms are shaped by selection to match features in the environment	Causation is a reciprocal process in which organisms shape and are shaped by selective developmental environments; phenotypic changes and acquired characteristics may lead rather than invariably follow genetic change
Directionality	Variation arises through random genetic mutations with no directionality	Variation arises through combination of genetic and constructive processes; biases in phenotypic variation provide directionality
Targets of Selection	Targets of selection are genes and alleles of genes; evolution consists of changes in gene frequencies	Targets of selection may be alleles, genes, organisms, or groups of organisms; evolution consists of change on any of these levels
Inheritance	Genetic inheritance is the only inheritance system; only genetically encoded traits can be inherited	Multiple systems shape transgenerational inheritance, both internal to the organism (genetic, epigenetic, maternal) and external (ecological, social learning, cultural); acquired traits may be inherited
Tempo & Pace of Change	Evolution proceeds at a gradual pace made up of micro-evolutionary processes shaped primarily by selection, but also drift, mutation and gene flow	Evolution proceeds at an uneven pace with periods of stasis punctuated by periods of rapid macro-evolutionary change

Table 2: Test Implications and data sets of explanatory frameworks for initial domestication

Category	Variables	Selectionist/ OFT Predictions	NCT Predictions	Archaeological Data
Context of initial domestication	Climate/Environmental Change	Change resulting in reduction of relative bio-mass	Either stable or improving, resulting in increase in relative bio-mass	Paleo-environmental proxies
	Population Levels	Marked increase or stable; reaching or surpassing carrying capacity	Little or no change; population levels remains well below carrying capacity	Number and size of sites and intensity of occupation
	Mobility	Decrease in mobility due to population packing	Decrease in mobility with settlements targeting resource-rich areas within circumscribed catchment, without population packing	Settlement pattern data and correlation with regional resource mapping
	Resource Depression	Decrease in high ranking resources. Intensification of procurement and processing activities	Broadening of resource base, with either no reduction in available biomass or intensification in processing	Archaeological Plant and Animal Remains
Role of niche construction in initial domestication	Modification of environments and biotic communities	Only adopted as a response to decline in availability in high ranking prey	Continuous and intensifying alteration of landscape or modification of biotic communities in absence of evidence of resource pressure	Evidence of human manipulation of environments (burning, changes in weed assemblages, appearance of species outside

				habitats)
Role of Social Behaviors	Social interaction, ritual behaviors, feasting	Intensification in these activities aimed at promoting competitive advantage of successful foragers over others	Intensification aimed at transmitting environmental knowledge, enhancing community cohesion, and leveling intra-community differences in access to resources.	Evidence of feasting, changes in community structure, burial practices
Pace of change	Dates of appearance of domesticates and agricultural emergence	Gradual and incremental	Periods of stasis and periods of rapid change in domesticates and humans	AMS Dating of material culture and archaeological materials, aDNA



Supplemental Information

Contrasting Neo-Darwinian and NCT informed explanatory frameworks

Table 2 in the main text contrasts distinct predictions about initial domestication grounded in neo-Darwinian and NCT explanatory frameworks. It also lists the kinds of archaeological data needed to test these predictions. Systematic head-to-head evaluations of the different neo-Darwinian and NCT predictions have yet to be performed in many world areas in which plants and animals were domesticated (figure 1). However, a review of empirical data from regions with the more complete records support NCT based predictions about the context of initial domestication, the role of niche constructing activities, the importance of social and ritual activities, and the pace of initial domestication and subsequent agricultural emergence.

The context of initial domestication

Neo-Darwinian explanatory frameworks predict that initial domestication universally occurred within the context of an imbalance between human population levels and environmental carrying capacity [1-3]. In contrast, NCT-based explanations predict that initial domestication is more likely to occur within environments in which abundant resources can be reliably found within circumscribed areas with population levels well below carrying capacity [4-8]. Evidence from multiple world areas supports NCT predictions on the context of initial domestication.

Arguments that initial domestication of South American crop plants was a direct response to the loss of high ranking game and plant resources [1], for example, are undermined by the 3000 years that separate the extirpation of Pleistocene mega-fauna and the appearance of humans in northern South America, and the subsequent 2000 year gap between human arrival and the earliest evidence of domesticates [7]. The stability of forest composition in regions where domesticates first appear in the Neotropics [9] also challenges the hypothesis that initial domestication was a response to a dramatic environmentally-driven change in resource availability.

In the Early Holocene Near East where ameliorating climatic conditions caused a rebound of biotic communities, neo-Darwinian models propose that rapid increases in human

population and subsequent population packing quickly outstripped the capacity of these resource-rich environments to support them, leading to an abandonment of more optimal mobile foraging strategies, resource diversification, intensification, and, subsequently, domestication [2]. These arguments, however, are based on a tenuous tautology that interprets evidence of loss of mobility as proof of a proposed causal linkage between population packing and loss of mobility [8]. Similar circularity is evident when it is claimed that apparent diversification of dietary resources provides support for the purported causal role of resource pressure in diversification. Perhaps more salient, there is no evidence of an imbalance between population and carrying capacity of other resource rich portions of the central and eastern Fertile Crescent where all four major livestock species (sheep, goat, cattle, and pig) and several crop plants were domesticated [10,11].

Recently Weitzl and Coddling have argued that a slight uptick in the number of radiocarbon dates from sites in eastern North America coincident with initial domestication of indigenous seed crops lends support to their thesis that population increase, and subsequent demographically-driven resource depletion, played a causal role in domestication in this region [3]. The Mid-Holocene was marked by a dramatic increase in resource availability in this region [6]; so it is not surprising to see some evidence of population increase at this time. But, as Weitzl and Coddling admit, there is no evidence for an imbalance between population and environmental carrying capacity at this time. Indeed, there is a good deal of evidence that initial domestication in eastern North America arose within resource rich environments in the absence any pressure on resource availability [12], leading even proponents of neo-Darwinian explanatory frameworks to admit that the example of initial domestication in eastern North America fails to “show a good fit” with predictions generated by models grounded in neo-Darwinian principles [1:140,13].

Instead, in regions where reasonably good settlement and paleo-environmental data exist (the Near East, eastern North America, and most recently China), initial domestication occurred in the absence of any demonstrable resource pressure, with population levels remaining well below likely carrying capacity. In each case the domestication process began in the context of small, semi- to fully-sedentary settlements strategically situated at the junction

of multiple resource rich eco-zones where inhabitants could draw on diverse and seasonally predictable resources drawn from circumscribed catchment areas capable of supporting these communities over extended periods of time [4,8-14] – conditions that closely conform to NCT predictions about the context of initial domestication specified in Table 2.

The role of niche construction in initial domestication

Neo-Darwinian models predict that niche-constructing activities leading to domestication are only adopted as a way of compensating for the lower returns of less intensive resource extraction strategies and are considered as sign of a decrease in foraging efficiency [3,15]. NCT frameworks, in contrast, predict that the practice of niche-constructing activities should not be seen as an adaptive response to changes in selective pressures in external environments (i.e., resource depression), but should instead be considered a means by which foragers actively manipulated selective pressures to their own evolutionary advantage. Thus, rather than being adopted late in the process of domestication as a last resort measure to combat declining returns from the environment, a NCT perspective predicts that these activities should be evident, in the absence of resource depression, prior to, or at the latest during, the initial stages of the domestication, and should intensify through time as a means of keeping carrying capacity in balance with population [5,6].

Once again emerging empirical evidence support NCT predictions. In northern South America, the Near East, eastern North America, and in China there is increasing evidence that foragers engaged in forest management through burning and coppicing, moved and tended and tending economically important plants, and harvested prey animals in ways that increase returns, in the absence of resource pressure, well before the appearance of archaeologically detectable domesticates [4-9,14,16-22].

The role of social and ritual activities

Social and ritual activities, if mentioned at all in explanatory frameworks grounded in neo-Darwinian principles, are usually portrayed as responses to tensions arising from resource depression and are directed toward increasing the competitive (and reproductive) advantage of

some foragers over others [23,24]. Under the NCT framework, in contrast, these activities are predicted to play key, catalytic roles as vehicles for transmitting environmental knowledge from one generation to the next, enhancing community cohesion, and combatting forces that encourage intra-group differences in access to basic resources [5].

The archaeological record of social and ritual activities needed to evaluate these predictions is sparse in many world areas. In the Near East, however, where there is a clear and steady uptick in social and ritual activities in Early Holocene communities on the threshold of domestication, NCT predictions find more support than those stemming from a Neo-Darwinian perspective. It is true that some researchers have interpreted the increase in ritual feasting in Early Holocene forager communities as evidence for increased competition, seeming to align more closely with neo-Darwinian predictions [24,25]. The absence of any evidence of status or economic differentiation between members of these small communities or signs that feasting was anything other than a broadly communal affair, however, have led the balance of researchers working in this area to conclude that these activities served an integrative role in combating potentials for inequalities likely to arise from increased investment in the manipulation and extraction of resources from more narrowly defined catchments areas [see multiple chapters in ref. 26, also 27,28].

The pace of initial domestication

Finally, under neo-Darwinian models the co-evolutionary relationships between humans and target domesticates at the heart of the process of domestication are predicted to unfold at a gradual, incremental pace as the result of the accumulation of thousands of decisions by individual foragers shaped by natural selection [29,30]. A NCT informed perspective, on the other hand, predicts that the process of domestication proceeds at an uneven pace, with periods of stasis punctuated by periods of rapid change, driven by both the rapid expression and accumulation of variation resulting from developmental processes that is turbo-charged by the human capacity for inventing and transmitting goal-directed behaviors [5].

Once again, the Near East provides us the most robust data sets regarding the pace of change in domestication and agricultural origins, and, once again, NCT predictions find more

empirical support. We now know that the record of initial domestication and subsequent agricultural origins in the Near East spans a period of about 2000 years – from about 11,700-10,000 years ago. Developments over this long period did not unfold as a series of gradual micro-evolutionary changes, however [31]. Instead the process was marked by a long period of relative stasis as small communities engaged in an extended process of broad spectrum resource extraction and increasing engagement in niche modification (from about 11,700-10,500), punctuated at about ca. 10,500 by a period of more rapid change as multiple different managed and morphologically domesticated crops and animals appear in different parts of the Fertile Crescent. This is followed by a several hundred year period in which domesticates play a relatively small role in the diet, interrupted by an additional period of rapid change which sees a virtual explosion of more fully committed agro-pastoral village communities across and out of the Fertile Crescent at 10,000-9,000 years ago [31-33].

References

- 1 Piperno DR. 2006 The origins of plant cultivation and domestication in the Neotropics: a behavioral ecological approach. In *Behavioral ecology and the transition to agriculture* (eds DJ Kennett, B Winterhalder), pp. 137–166. Berkeley, CA: University of California Press.
- 2 Stutz AJ, Munro ND, Bar-Oz G. 2009 On increasing the resolution of the Broad Spectrum Revolution in the southern Levantine Epipaleolithic (19–12 ka). *J Hum Evol* 56, 294–306.
- 3 Weitzel EM, Coddling BF 2017 Population growth as a driver of initial domestication in Eastern North America. *Royal Society Open Science*.
- 4 Crawford G. 2011 Advances in understanding early agriculture in Japan. *Curr Anthropol* 52, S331–S345.
- 5 Zeder MA. 2016 Domestication as a model system for niche construction theory. *Evol Ecol* 30,325-348.
- 6 Zeder MA, Smith BD. 2009 A conversation on agriculture: talking past each other in a crowded room. *Curr Anthropol* 50, 681–691.

- 7 Smith BD. 2016 Neo-Darwinism, niche construction theory, and the initial domestication of plants and animals. *Evol Ecol* 30, 307-324.
- 8 Zeder MA. 2012a The broad spectrum revolution at 40: resource diversity, intensification, and an alternative to optimal foraging explanations. *J Anthropol Archaeol* 31, 241–264.
- 9 Gnecco, C. (2003). Against ecological reductionism: Late Pleistocene hunter-gatherers in the tropical forests of northern South America. *Quatern Int* 109–110, 13–21.
- 10 Zeder MA. 2011. The origins of agriculture in the Near East. *Curr. Anthropol.* 54 (S4), S221-S235.
- 11 Arbuckle BS 2015. Large game depression and the process of animal domestication in the Near East. In: *Climate Change in Ancient Societies*, (eds MS Kerner, R Dann, P Bangsgaard), pp. 215-43 . Chicago: University of Chicago Press.
- 12 Smith BD, Yarnell RA. 2009. Initial formation of an indigenous crop complex in eastern North America at 3800 B.P. *Proc Natl Acad Sci USA* 106(16), 6561-6566.
- 13 Gremillion K. 2004 Seed processing and the origins of food production in eastern North America. *Amer Antiq* 69, 215–233.
- 14 Ren X, Lemoine X, Mo D, Kiffer TR, Guo Y, Qin Z, Liu X, 2016 Foothills and intermountain basins: does China's fertile arch have 'Hilly Flanks? *Quat Int* 334+335, 165-178.
- 15 Munro ND, Atici, L. 2009 Human subsistence change in the Late Pleistocene Mediterranean Basin: the status of research on faunal intensification, diversification, and specialization. *Before Farming: The Archaeology of Old World Hunter–Gatherers* 2009/1, article 1.
- 16 Piperno, D, Bush M, Colinvaux P. 1991a). Paleoecological perspectives on human adaptation in Panama I: The Pleistocene. *Geoarchaeology* 6, 201–226.
- 17 Piperno D, Bush M, Colinvaux P. 1991a. Paleoecological perspectives on human adaptation in Panama I: The Pleistocene. *Geoarchaeology* 6, 201–226.

- 18 Aceituno FJ, Loaiza N, Delgado-Burbano M, Barrientos G. 2013 The initial settlement of Northwest South America during the Pleistocene–Holocene transition: synthesis and perspectives. *Quatern Int* 30, 23–33.
- 19 Asouti E, Kabukcu C, White CE. 2015. Early Holocene woodland vegetation and human impacts in the arid zone of the southern Levant. *Holocene*. 25, 1565-1580.
- 20 Delcourt P, Delcourt H, Ison C, Sharp W, Gremillion K. 1998 Prehistoric human use of fire, the eastern agricultural complex, and Appalachian Oak-Chestnut forests: Paleoecology of Cliff Palace Pond, Kentucky. *Antiq* 6,12,14-20].63, 263–278.
- 21 Gremillion K. 1998 Changing roles of wild and cultivated plant resources among early farmers of eastern Kentucky. *Southeast Archaeology* 17, 140–157.
- 22 Watling J, Iriarte J, Mayle FE, Schaan D, Pessenda LCR, Loader NJ, Street-Perrott FA, Dikau RE, Damasceno A, Ranzi A. 2017 Impact of pre-Columbian "geoglyph" builders on Amazonian forests. *PNAS USA* 114:1868-1873.
- 23 Aldenderfer, M., 2006. Costly signaling, the sexual division of labor, and animal domestication in the Andean Highlands. In *Behavioral Ecology and the Transition to Agriculture*. eds DJ Kennett, B Winterhalder, pp. 167–196. Berkeley, CA: University of California Press.
- 24 Starkovich BM., Stiner MC. 2009 Hallan Çemi Tepesi: high ranked game exploitation alongside intensive seed processing at the Epipaleolithic Neolithic transition in southeastern Turkey. *Anthropozoologica* 44, 41–61.
- 25 Hayden B. 1995 A new overview of domestication. In: *Last Hunters, First Farmers: New Perspectives on the Transition to Agriculture*, eds TD Price, A-B Gebauer, pp. 273-300. Santa Fe, NM: School of American Research Press.
- 26 Kuijt I. ed 2000 *Life in Neolithic farming communities: social organization, identity, and differentiatio*. New York, NY: Plenum.
- 27 Zeder MA. 2012c Religion and the revolution: the legacy of Jacques Cauvin. *Paléorient* 37(1), 39–60.

- 28 Zeder MA, Spitzer MD. 2016. New insights into broad spectrum communities of the Early Holocene Near East: the birds of Hallan Çemi. *Quat Sci Rev* 151, 140-159.
- 29 Rindos D. 1984 *The origins of agriculture: an evolutionary perspective*. Orlando, FL: Academic Press.
- 30 Winterhalder B, Kennett DJ (2006) Behavioral ecology and the transition from hunting and gathering to agriculture. In *Behavioral ecology and the transition to agriculture*, eds DJ Kennett, B Winterhalder, pp. 1–21. Berkeley, CA: University of California Press.
- 31 Zeder MA. 2009 Evolutionary biology and the emergence of agriculture: the value of co-opted models of evolution in the study of culture change. In: *Macroevolution in human prehistory* (eds A Prentiss, I Kuijt, J Chatters), pp. 157–210. New York, NY: Springer.
- 32 Kuijt I, Prentiss AM. 2009 Niche construction, macroevolution, and the Late Epipaleolithic of the Near East. In *Macroevolution in human prehistory* (eds AM Prentiss, I Kuijt, J Chatters), pp. 253–271. New York, NY: Springer.
- 33 Byrd B. 2005. Reassessing the emergence of village life in the Near East. *J Archaeol Res* 13, 231–290.